

Structure of the La motif: a winged helix domain mediates RNA binding via a conserved aromatic patch

Gang Dong¹, Ghadiyaram Chakshumathi¹, Sandra L Wolin^{1,2} and Karin M Reinisch^{1,*}

¹Department of Cell Biology, Yale University School of Medicine, New Haven, CT, USA and ²Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, CT, USA

The La protein is a ubiquitous nuclear phosphoprotein that recognizes the 3' uridylates found in all newly synthesized RNA polymerase III transcripts. La binding stabilizes these transcripts from exonucleases and may also assist their folding. Here we present the first structural insights into how the La protein specifically interacts with its RNA substrates. The most conserved region of the La protein is the La motif, a domain also found in several other RNA-binding proteins. We have determined the structure of the La motif from the *Trypanosoma brucei* La protein to 1.6 Å resolution (PDB code 1S29). The La motif adopts a winged helix–turn–helix architecture that has a highly conserved patch of mainly aromatic surface residues. Mutagenesis experiments support a critical role for this patch in RNA binding and show that it partly determines binding specificity for RNAs ending in 3' hydroxyl, a defining characteristic of the La protein. These findings reveal that the La motif is essential for high-affinity binding and also contributes to specificity.

The EMBO Journal (2004) 23, 1000–1007. doi:10.1038/sj.emboj.7600115; Published online 19 February 2004

Subject Categories: structural biology; RNA

Keywords: La autoantigen; RNA-binding protein; RNA–protein interactions; tRNA processing; winged helix domain

Introduction

The La protein was first described as an autoantigen in patients with the rheumatic diseases systemic lupus erythematosus and Sjogren's syndrome (Mattioli and Reichlin, 1974; Alspaugh and Tan, 1975). Although first characterized in humans, La homologs have since been found in all eucaryotes including trypanosomes, yeast, and plants (see Maraia and Intine, 2001; Wolin and Cedervall, 2002).

La proteins from humans, *Drosophila*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Trypanosoma brucei* have been characterized (Chambers *et al.*, 1988; Bai *et al.*, 1994; Yoo and Wolin, 1994; Van Horn *et al.*, 1997; Marchetti *et al.*, 2000; Wolin and Cedervall, 2002). La is

localized primarily in the nucleus, where it binds nascent RNA polymerase III transcripts. RNAs bound by La include pre-tRNAs, 5S rRNAs, U6 RNA, and SRP RNA (see Wolin and Cedervall, 2002). In all of these RNAs, La specifically recognizes the 3' UUU_{OH} as part of its binding site (Stefano, 1984; Wolin and Cedervall, 2002). In the yeast *S. cerevisiae*, La also binds nascent RNA polymerase II transcripts such as U1, U2, U3, U4, and U5 snRNA that have been partially processed to end in polyuridine (Kufel *et al.*, 2000; Xue *et al.*, 2000). One characteristic feature of La is its preference for a hydroxyl over a phosphate group at the 3' terminus. This may allow La to distinguish intact nascent polymerase III transcripts from degraded RNA molecules with 3' terminal phosphates and to discriminate between 3' terminal and internal polyuridine stretches (Stefano, 1984; Long *et al.*, 2001). So far, the best-established role for La is to protect newly synthesized RNAs from exonucleases, with various consequences for different transcripts. For pre-tRNAs, La is required for the normal endonucleolytic removal of the 3' trailer (Yoo and Wolin, 1997) and may also be important for their 5' maturation (Fan *et al.*, 1998). For certain pre-tRNAs, binding by La facilitates correct folding (Chakshumathi *et al.*, 2003). For other transcripts, La binding facilitates their assembly into functional ribonucleoprotein particles (Pannone *et al.*, 1998; Xue *et al.*, 2000). Although La has also been proposed to be a transcription factor for RNA polymerase III (Gottlieb and Steitz, 1989; Maraia, 1996), it is not required for transcription in human, *Xenopus*, or yeast cell extracts (Yoo and Wolin, 1997; Lin-Marq and Clarkson, 1998; Chong *et al.*, 2001).

Like many RNA-binding proteins, La is modular. An N-terminal 'La motif' (LM) is followed by either one or two RNA recognition motifs (RRMs, also referred to as RBDs or RNP domains) (see Varani and Nagai, 1998; Wolin and Cedervall, 2002). The size of La varies from ~32 kDa in yeasts, where La has only one RRM, to 50 kDa in humans, where La has two RRM. The most C-terminal region of La is highly charged and is known to be phosphorylated (Fan *et al.*, 1997; Long *et al.*, 2001). Whereas sequence identity breaks down in the RRM and C-terminal regions of La, the LM is conserved across species. Human La, for example, has 29% sequence identity with the *S. cerevisiae* and 40% identity with the *T. brucei* N-terminal domain. In all characterized La proteins, the LM is at the N-terminus (Wolin and Cedervall, 2002).

The LM is also found in a number of other proteins that are otherwise structurally unrelated to La. Two of these proteins, the closely related *S. cerevisiae* proteins Slf1p and Sro9p, are polyribosome associated and may function in translation (Sobel and Wolin, 1999). The other characterized LM protein, p43 from the ciliate *Euplotes aediculatus*, is a component of telomerase (Aigner *et al.*, 2003). In contrast to the La protein, p43 lacks sequence specificity for the 3' UUU_{OH} (Aigner *et al.*, 2003).

Both the LM and the central RRM are necessary for polyuridine specific binding (Goodier *et al.*, 1997; Jacks *et al.*, 2003).

*Correspondence. Department of Cell Biology, Sterling Hall of Medicine, Yale University School of Medicine, 333 Cedar St, New Haven, CT 06520-8002, USA. Tel.: +1 203 785 6469; Fax: +1 203 785 7446; E-mail: karin.reinisch@yale.edu

Received: 21 November 2003; accepted: 16 January 2004; published online: 19 February 2004

The role of the second RRM that is present in some species is not understood. It does not appear to contribute to binding polyU (Goodier *et al*, 1997; Ohndorf *et al*, 2001; Jacks *et al*, 2003). Moreover, in the solution structure of this domain (Jacks *et al*, 2003), the canonical RNA-binding surface is acidic, suggesting that it may not interact with RNA at all. Possibly then, this domain is a module for interacting with other proteins.

Here we begin to address how the La protein specifically interacts with its RNA targets. We present the 1.6 Å crystal structure of the LM domain from the *T. brucei* La protein. The LM adopts a winged helix–turn–helix (wHTH) fold present in many transcription regulatory proteins (Gajiwala and Burley, 2000). An aromatic patch on the domain surface is strictly conserved in all LMs, and structure-based mutagenesis studies show that residues in this patch are critical for high-affinity RNA binding and 3' end recognition.

Results and discussion

Overview of the structure: the La motif is a variation of the winged helix–turn–helix fold

To obtain crystals of the La protein, we screened constructs based on the *T. brucei*, *S. pombe*, *S. cerevisiae*, and human La proteins. Crystals were obtained for the LM domain from *T. brucei*. The LM construct used consists of residues 1–89 of La and an additional Gly-Ser-His at the N-terminus. Although these three residues are not part of the *T. brucei* sequence, they are essential in forming crystal contacts. The structure with 100 water molecules has been refined to 1.6 Å with $R = 23.2\%$ / $R_{\text{free}} = 25.0\%$ and excellent geometry (Table I).

Table I Summary of crystallographic data and refinement

<i>Data collection</i>	
Space group	P2 ₁ 2 ₁ 2 ₁
Cell constants (Å)	$a = 32.926$, $b = 53.833$, $c = 53.912$
Wavelength (Å)	0.9796
Resolution (Å)	1.6
Last shell (Å)	1.65–1.59
No. of unique reflections (last shell)	24 672 (2382)
Redundancy (last shell)	7.0 (4.1)
R_{sym} (last shell) (%)	3.2 (6.0)
I/σ (last shell)	41 (21)
Completeness (last shell) (%)	99.5 (95.7)
Figure of merit (last shell)	0.45 (0.40)
Phasing power (last shell)	2.52 (1.78)
<i>Refinement</i>	
Resolution (Å)	20.0–1.6
$R_{\text{work}}/R_{\text{free}}$ (%)	23.2 (25.0)
rmsd bond lengths (Å)	0.00526
rmsd bond angles (deg)	1.169
rmsd improper angles (deg)	0.76
rmsd dihedral angles (deg)	21.0
Residues in favored Ramachandran region	91
Residues in additionally allowed region	1
Residues in generously or disallowed region	0
B value rmsd (main/side)	1.0/2.2
No. of water molecules	100

$R_{\text{sym}} = \frac{\sum_j \sum_i |I_j - \langle I \rangle|}{\sum_j I_j}$
Phasing power = $\frac{\sum |F_+ - F_-|^2}{\sum |E|^2}^{1/2}$, where the denominator is the lack of closure error.

$R_{\text{work, free}} = \frac{\sum (|F_{\text{obs}}| - |F_{\text{calc}}|)}{\sum |F_{\text{obs}}|}$ for the reflections used in refinement (work) and 10% of reflections held aside (free).

The LM domain is mostly α -helical, consisting of six helices and a three-stranded antiparallel sheet (Figure 1A). One long α -helix (H1) and two shorter ones (H1' and H2) precede the first β -strand (B1). Helices H3, H4, and H5, arranged to form a 'C' shape, reside between β -strands B1 and B2, and a six-residue loop links B2 and B3. Overall, the LM adopts a wHTH fold found in both eucaryotic and prokaryotic transcription factors (Figure 1B) (Gajiwala and Burley, 2000). Two of the three N-terminal helices and H4 in the LM may be viewed as insertions into the wHTH topology. The wHTH transcription factors typically contain a three-helix bundle and a three-stranded antiparallel β -sheet. H1 precedes B1; H2 and H3 reside between B1 and B2; and a loop, the wing, links B2 and B3. In some wHTH proteins, a second wing follows B3. Winged HTH domains have different functions. In *FokI* restriction endonuclease and replication protein A, for example, the wHTH is involved in protein–protein interactions (Wah *et al*, 1997; Mer *et al*, 2000). In many transcriptional regulators, it is involved in binding dsDNA. Most frequently, wHTHs interact with dsDNA by inserting the recognition helix H3 (H5 in the LM) into the DNA major groove, and often the wing also makes contacts with the DNA (Gajiwala and Burley, 2000).

To our knowledge, the elongation factor selB is the only previously known wHTH protein that binds RNA. The C-terminus of selB consists of four consecutive wHTH domains, where the two most C-terminal of these interact with an mRNA hairpin (Fourmy *et al*, 2002; Selmer and Su, 2002). The residues that bind RNA have not been identified. Sequence conservation and electrostatic potential calculations point to residues in the most C-terminal wHTH located on a surface between H3 (H5 in La), H2 (H3 in La), and the β -hairpin between B2 and B3 (Selmer and Su, 2002). This surface is similar to that used for DNA recognition in many of the wHTH transcription factors. In La, the residues on the corresponding surface are not well conserved. As discussed below, the LM may use an entirely different, highly conserved surface between helices H1, H1', H2 and the loop between H3 and H4 to interact with RNA (Figures 1A and 2).

Conserved residues form an aromatic patch on the surface of La

LM-containing proteins share the highest sequence identity with the more N-terminal portions of the *T. brucei* LM




Figure 1 (A) Ribbon views of the LM of *T. brucei* La. (B) Ribbon view of a wHTH motif as observed in the Z-DNA-binding domain of the adenosine deaminase ADAR1 (PDB accession number 1QBJ). The topology of the LM is similar to that of the wHTH motif of some transcription factors. MOLSCRIPT (Kraulis, 1991) was used to prepare this figure.




Figure 3 Binding of wild-type and mutant La proteins to RNA. (A) ³²P-labeled yeast pre-tRNA^{Arg}_{CCG} (left panel) or pre-tRNA^{Arg}_{CCG} lacking the 9 nt 3' trailer (right panel) was incubated without protein (lane 1) or with the truncated *T. brucei* La protein (LM+RRM; amino acids 1–192) at the indicated protein concentrations. The pre-tRNA concentration in the reaction was 0.15 nM. Naked RNAs and RNPs were separated in native gels. (B) ³²P-labeled yeast pre-tRNA^{Arg}_{CCG} (final concentration 0.15 nM) was incubated with either no protein (lane 1) or the indicated concentrations of full-length La (lanes 2–4), La 1–192 (lanes 5–8), the isolated LM (amino acids 1–89; lanes 9–11), the RRM (residues 94–192; lanes 12–14), or a 1:1 mixture of the LM and RRM (lanes 15–17). Naked RNAs and RNPs were separated by native gel electrophoresis. (C) Binding of La 1–192 (left panel) and the D27A mutant (right panel) to pre-tRNA^{Arg}_{CCG} was performed as described in (A). Protein concentrations in the reactions are indicated above the lanes. In lanes 1 and 10, no protein was added. We quantitated the difference in binding affinity of La 1–192 and the D27A mutant by performing multiple experiments. These revealed that La 1–192 bound the RNA with a K_d of 4.1 ± 0.7 nM, while the D27A mutant bound the RNA with a K_d of 14.6 ± 2.9 nM, an approximately 3.6-fold decrease in affinity. (D) pre-tRNA^{Arg}_{CCG} was subjected to gel shift analysis using either the wild-type La 1–192 (lanes 2–4) or the mutants Q14A/E16A (lanes 5–7), F29A (lanes 8–10), F50A (lanes 11–13), F17A (lanes 14–16), and Y18A (lanes 17–19). Protein concentrations in each lane are given in nM. Lanes 1 and 20, no protein.

shown). In the experiment shown, approximately 50% of the labeled pre-tRNA was shifted into RNA–protein complexes at a protein concentration between 4.8 and 9.6 nM. Deleting the

UUU_{OH}-containing trailer or replacing it with polycytosine reduced La binding significantly (Figure 3A, right panel, and data not shown).

As with human La (Goodier *et al*, 1997; Ohndorf *et al*, 2001; Jacks *et al*, 2003), both the *T. brucei* LM and the adjacent RRM are necessary for high-affinity RNA binding. Neither the LM (residues 1–89) nor the RRM (residues 94–192) alone bind to RNA with high affinity (Figure 3B, lanes 9–11 and 12–14), but a construct consisting of both the LM and the adjacent RRM (residues 1–192) binds pre-tRNA with a similar affinity to the full-length protein (compare lanes 2–4 with lanes 5–7). Interestingly, a 1:1 mixture of the LM and the RRM also fails to bind pre-tRNA (Figure 3B, lanes 15–17). As the linker region between the two domains (residues ~89–96 in *T. brucei*) varies in length and is not well conserved in La proteins from different species, it seems unlikely that it is involved in important protein–RNA contacts. Most likely, the LM and the RRM do not have an extensive protein–protein interface and are flexibly tethered in the absence of RNA. Nucleic acid binding may orient the domains as is the case for Sxl, which binds ssRNA via two adjacent RRMs, or transcription factor TFIIIA, which binds dsDNA with three tandem zinc fingers (Wuttke *et al*, 1997; Crowder *et al*, 1999).

Residues in the conserved aromatic patch are essential for La interactions with pre-tRNA

To test which residues in the LM are important for RNA binding, we constructed a series of point mutations in the LM+RRM construct (residues 1–192). We mutated conserved surface residues in the LM to alanine and assayed the binding affinity for each construct by EMSA with pre-tRNA (Table II). Mutations of conserved surface residues not in the aromatic patch (the double mutations R83A/R84A and E39A/F41A) did not significantly affect RNA binding, as the amount of mutant protein required to shift half the labeled pre-tRNA varied less than two-fold from the wild type (data not shown). Mutation of three conserved residues in the patch (R52A, N23A, and D27A) all resulted in modest decreases in binding affinity compared to wild-type protein (shown for D27A in Figure 3C; Table II). In contrast, the double mutation Q14/E16A and mutations of aromatic residues in the conserved patch (F29A, F50A, F17A, Y18A, and the double mutation F17A/Y18A) diminish the RNA-binding affinity by more than 100-fold (Figure 3D). These results confirm that residues in the LM conserved patch are required for La interactions with RNA.

Table II Effect of mutations on RNA-binding affinity

Mutation	Location	Effect on RNA-binding affinity
Q14A/E16A	Conserved patch	> 100 × reduced
F17A	Conserved patch	> 100 × reduced
Y18A	Conserved patch	> 100 × reduced
F29A	Conserved patch	> 100 × reduced
F50A	Conserved patch	> 100 × reduced
N23A	Conserved patch	Slightly reduced (~2–4 ×)
D27A	Conserved patch	3.6 × reduced
R52A	Conserved patch	Slightly reduced (~2–4 ×)
R83A/R84A	Not in patch	< 2 × reduced
E39A/F41A	Not in patch	< 2 × reduced

Mutating an aspartate in the conserved surface patch reduces La specificity for a 3' hydroxyl versus phosphate group

One of the more interesting features of La recognition is its ability to distinguish RNAs ending in UUU_{OH} from those ending in UUU_p. We used competition experiments to examine those mutants that retained RNA-binding affinity (R52A, E39A/F41A, R83/R84, N23A, and D27A) for their ability to discriminate between RNAs differing only at the 3' ends. In these experiments, we determined the amount of unlabeled competitor RNA necessary to dissociate the labeled pre-tRNA complex. Pre-tRNA containing a 3' UUU_{OH} was a more effective competitor for *T. brucei* La binding than the same RNA terminating in UUU_p (Figure 4A), as previously described for the human and the *S. cerevisiae* La proteins (Stefano, 1984; Long *et al*, 2001). Mutations of residues outside the aromatic patch (E39A/F41A and R83A/R84A) did not affect the ability of *T. brucei* La to recognize 3' OH, nor did the R52A mutation at the edge of the aromatic patch (data not shown). The N23A mutation in the patch also had little, if any, effect (Figure 4B). In contrast, the D27A mutation at the base of the patch has a diminished preference for the hydroxyl group, as less UUU_p-containing pre-tRNA was needed to dissociate the complex than required for the wild-type protein (compare Figures 4C and A, left panels). For the wild-type protein, the amount of 3' UUU_{OH} competitor required to dissociate 50% of the labeled protein–RNA complex ($K_d^{1/2}$) was 9.9 ± 0.9 nM, while for the 3' UUU_p competitor it was 988.6 ± 70.3 nM (Figure 4D). For the D27A mutant, these values were 19.9 ± 1.7 and 358.3 ± 62.3 nM, respectively (Figure 4D). For the native protein the relative binding affinity ($K_{rel} = K_d^{1/2} \text{ UUU}_p / K_d^{1/2} \text{ UUU}_{OH}$) was approximately 100, while for the D27A mutant the ratio was approximately 18, a five- to six-fold decrease in specificity. D27 then contributes at least a part of the specificity of the *T. brucei* La protein for the 3' hydroxyl. In the native protein, D27 could hydrogen bond to the 3' OH while sterically and electrostatically clashing with a negatively charged 3' phosphate group. Replacing the aspartate with alanine would at least partially remove steric and electrostatic barriers that prevent a phosphate group from binding, thus resulting in diminished discrimination against it.

Models for RNA binding

The La protein makes specific interactions with the 3' UUU_{OH}, optimally accommodating four uridine bases in its recognition site (Stefano, 1984), and it may also interact with other regions of its cognate RNAs (Wolin and Cedervall, 2002; Chakshumathi *et al*, 2003). We have found that residues in the conserved aromatic patch of the LM are essential for high-affinity pre-tRNA binding. However, since RNA binding by La requires both the LM and the RRM, any model must take into account the role of the RRM. One possibility is that the LM functions indirectly to facilitate binding by the RRM. For example, the conserved aromatic patch may contribute to an LM:RRM interface that induces an RRM conformation able to interact with RNA. Our gel shift experiments show that the LM and RRM do not form a complex when separately expressed and then mixed together, either in the presence or absence of RNA (Figure 3B and data not shown), suggesting that the two domains do not share an extensive interface.




Figure 4 Mutating D27 reduces the specificity of La for RNAs ending in a 3' hydroxyl. (A) La 1–192 (4.8 nM) was mixed with ³²P-labeled pre-tRNA_{CCG}^{Arg} (0.15 nM) in the presence of a 0- to 10 000-fold molar excess of unlabeled pre-tRNA_{CCG}^{Arg} terminating in either UUU_p (left panel) or UUU_{OH} (right panel). Naked RNAs and RNPs were separated by native gel electrophoresis. The small differences in the fraction of RNA bound with 4.8 nM protein between this experiment and Figure 3C, lane 5, reflect experimental variation. Lane 1, no added protein. (B, C) The N23A mutant (B) or D27A mutant protein (C) was mixed with ³²P-labeled pre-tRNA_{CCG}^{Arg} (0.15 nM) in the presence of the indicated molar excess of unlabeled pre-tRNA_{CCG}^{Arg} as described in (A). The concentration of protein used in the reactions was 19.2 nM. Lane 1, no added protein. (D) Data from the competition titrations for La 1–192 and the D27A mutant are plotted as the fraction of labeled probe bound at each concentration of competitor RNA. Each data point represents the mean from at least three experiments. RNA concentrations are given in nM.

Nonetheless, we cannot exclude the possibility that the LM is only indirectly involved in RNA binding.

We favor a model in which the LM interacts directly with RNA. This is plausible in that the LM is the most conserved portion of La and is found in other RNA-binding proteins that lack recognizable RRM (Sobel and Wolin, 1999; Wolin and Cedervall, 2002). Our result, that mutating residue D27 in the LM conserved aromatic patch to alanine reduces La's specificity for a 3' OH over a 3' phosphate, supports a role for the conserved patch in forming at least part of the recognition site for 3' ends. With its many aromatic residues, the patch is well suited to interacting with one or more of the 3' uridine

bases. Since RRM is known to bind ssRNA and the La RRM is required for RNA binding, surfaces from both the LM and the RRM may contribute to the polyuridine-binding site. For both domains to bind to the relatively short RNA recognition sequence, they would have to form a V-shaped crevice lined on one side by the conserved aromatic patch of the LM and on the other by the RNA-binding surface of the RRM. The polyuridine would bind in this crevice, uridine bases interacting with each of the two domains. The length of the linker between the LM and the RRM varies in La proteins from different sources, and at eight amino acids the *T. brucei* linker is among the shortest. It is long enough, however, to allow juxtapositioning of the conserved aromatic La patch and the RNA-binding surface of the RRM. Since the LM and its conserved aromatic patch also occur in proteins that do not specifically bind UUU_{OH} (Sobel and Wolin, 1999; Aigner *et al*, 2003), it is possible that the specificity for polyU is a function of cooperative interactions by the LM and the RRM.

In summary, by combining structural information with mutational analysis, we have established that a conserved aromatic patch in the LM is required for high-affinity interactions of the La protein with RNA. Further, since D27 in the conserved aromatic patch contributes to the specificity for a hydroxyl versus a phosphate 3' terminus, the patch plays a role in binding the 3' terminus, possibly also contributing to the specificity of La for polyU. How the LM and the RRM cooperate in RNA binding and the precise nature of the protein-RNA interactions will best be resolved by a structure of the complex.

Materials and methods

Cloning

A full-length construct encoding *T. brucei* La was a generous gift of Dr Elisabetta Ullu. Wild-type full-length protein (residues 1–335) and three truncated versions (LM+RRM: residues 1–192; LM: residues 1–89; and central RRM: residues 94–192) of the *T. brucei* La protein were cloned into the plasmid pET-15b (Novagen) between the *NdeI* and *BamHI* restriction sites. Each construct includes a thrombin-cleavable hexahistidine tag at the protein N-terminus. As the LM+RRM construct has the same affinity for pre-tRNA as the full-length protein, it was used as a template for mutagenesis. A total of 11 mutations (Table II) were introduced into the LM+RRM construct using the QuickChange site-directed mutagenesis kit (Stratagene). All plasmids were sequenced to confirm the correct sequence and the presence of the expected mutation.

Protein overexpression and purification

Escherichia coli BL21(DE3) cells harboring plasmid for the full-length La, La+RRM, the LM, or the central RRM were grown at 37°C in Luria broth supplemented with 50 µg/ml ampicillin. When the cells reached an OD₆₀₀ between 0.6 and 0.8, they were shifted to 25°C (18°C for mutations) and protein production was induced with 1 mM IPTG. The cells were harvested 6–8 h after induction and stored at –80°C until further use. The protein purification was carried out entirely at 4°C. Frozen cells were thawed at room temperature and resuspended in 20 ml/l cell culture of cold lysis buffer (20 mM Tris-HCl (pH 8 at room temperature), 300 mM NaCl, 1 mM PMSF, and 0.5 mg/ml lysozyme). After a brief incubation, the cells were further lysed by pulse sonication. The lysate was cleared by centrifugation at 30 000 g for 30 min, followed by filtration through a 0.4 µm filter. The lysate was supplemented with 10 mM imidazole and loaded onto Ni-NTA resin (Qiagen) pre-equilibrated with buffer A (20 mM Tris-HCl (pH 8.0 at room temperature), 300 mM NaCl, and 10 mM imidazole). The resin-bound protein was washed with 10 volumes of buffer A. Thrombin (150 U/l of cell culture; Amersham) was added to release the protein from the resin. Protein, consisting of an N-terminal Gly-Ser-His sequence followed by the desired *T. brucei* La sequence, was washed from the column

with 5–7 bed volumes of buffer A. The protein was concentrated to a small volume in the presence of 10% glycerol and loaded onto a Superdex200 gel filtration column (Amersham). Protein was collected from the gel filtration column in buffer B [10 mM Tris-HCl (pH 8.0 at room temperature), 100 mM NaCl, 5% v/v glycerol, and 5 mM MgCl₂]. LM+RRM mutants behaved like the native construct on the gel filtration column. Fractions corresponding to La were pooled and concentrated to 10–20 mg/ml, suitable for crystallization.

Selenomethionine-substituted LM was expressed using minimal media supplemented with all amino acids (2 mg/ml) except methionine. Just prior to induction, L-selenomethionine was added to 100 mg/l, and additional threonine, lysine, phenylalanine, leucine, isoleucine, and valine were added to inhibit the methionine biosynthetic pathway (Doublet, 1997). The selenomethionine protein was purified as described above, except for the addition of extra reductant. All steps preceding the gel filtration column were carried out in the presence of 20 mM β-me; after the Ni-NTA affinity purification, the protein was stabilized by the presence of 10 mM DTT.

Crystallization

Crystals of the LM were grown at 4°C by the hanging drop method against a well liquor containing 100 mM HEPES (pH 7.5), 5% (v/v) glycerol, 32–36% (w/v) PEG 3.5K, and, in the case of the selenomethionine-substituted protein, 10 mM DTT. Drops contained 1–6 µl of protein solution and an equivalent volume of well liquor. Crystals appeared within 1–2 days and grew to ~0.2 × 0.2 × 0.8 mm within a week. The crystals belong to space group P2₁2₁2₁ (*a* = 32.926 Å, *b* = 53.833 Å, and *c* = 53.912 Å) and contain one molecule per asymmetric unit. We were unable to crystallize the LM domain lacking the N-terminal Gly-Ser-His addition. The crystals were soaked in well liquor for about half an hour and flash-frozen in liquid nitrogen (Rodgers, 1997) prior to data collection. They diffracted to ~1.9 Å resolution in-house and to beyond 1.6 Å resolution at beamline X12C at Brookhaven National Laboratory and had a mosaicity of ~0.5°. Data from selenomethionine-substituted crystals were not isomorphous with crystals of the unsubstituted protein although both crystal forms belonged to the same space group and had the same unit cell dimensions.

Structure determination

Data from a flash-frozen selenomethionine-substituted crystal were collected at the beamline X12C at Brookhaven National Laboratory. Full and highly redundant data sets collected at the anomalous peak wavelength sufficed to calculate high-quality electron density maps (Table I). Data were integrated and scaled using the program HKL2000 (Otwinowski and Minor, 1997). All subsequent calculations were performed using the software suite CNS (Brunger *et al*, 1998). Anomalous Patterson maps allowed us to identify three selenium sites. These were refined in CNS, and SAD phases together with solvent flipping produced highly interpretable experimental maps at 1.6 Å. The initial model was built in O (Kleywegt and Jones, 1997). Subsequent cycles of torsion angle dynamics, least squares minimization, individual B-factor refinement, and the addition of water molecules yielded *R*_{work} = 23.2% and *R*_{free} = 25.0% for all data between 20 and 1.6 Å. The final model contains all residues including Gly-Ser-His (–3, –2, –1) and 1–89 of the *T. brucei* La sequence and 100 water molecules. Coordinates and structure factors have been deposited in the Protein Data Bank (accession number 1S29).

Electrophoretic mobility shift assays

The clone containing pre-tRNA^{Arg}_{CCG} under control of the T7 promoter has been described (Chakshusmathi *et al*, 2003). After linearization with *DraI*, transcription with T7 RNA polymerase yielded a pre-tRNA containing three U's at the 3' end. pre-tRNA^{Arg}_{CCG} sequences lacking the 3' trailer were amplified from the full-length clone using 5'-CGGCGAATTCTAATACGACTACTATAGTTTATTATGCTCCTCTAG TGC-3' and 5'-GCTCCTCCCGGACTCGAACC-3' and transcribed using T7 RNA polymerase. Pre-tRNAs were synthesized as described (Yisraeli and Melton, 1989), except that 50 µCi of [α-³²P]rCTP (400 Ci/mmol; Amersham) was used in place of rCTP. RNAs were gel purified and quantitated by scintillation counting. Prior to their use in binding assays, RNAs were resuspended in 50 mM Tris (pH 8.0) and 10 mM MgCl₂, refolded by heating to 60°C

for 1 min, and then slowly cooled to room temperature. For binding assays, 1.5 fmol of RNA was mixed with 6–768 fmol of protein and 0.2 µg/ml *E. coli* tRNA in binding buffer (20 mM Tris, pH 8.0, 3 mM MgCl₂, 0.1 mM EDTA, 2 mM DTT, 100 mM NaCl, and 0.1% NP-40). The total reaction volume was 10 µl. Reactions were incubated at room temperature for 10 min and then on ice for 10 min, and loaded onto 5% polyacrylamide (38:2 acrylamide:bisacrylamide)/5% glycerol gels (prerun at 16 V/cm for 10 min at 4°C). Gels were run at 16 V/cm in 0.5 × TBE (50 mM Tris, 45 mM boric acid, and 1.25 mM EDTA) at 4°C for 2 h and 30 min. The fraction of bound RNA (θ) was quantitated using a PhosphorImager (Molecular Dynamics). Using Imagequant, the data were fitted by nonlinear least squares analysis as a function of protein concentration using the equation

$$\theta = \frac{[\text{protein}]}{[\text{protein}] + K_d}$$

where θ is the fraction of bound RNA and K_d is the dissociation constant. Each K_d was calculated from at least three binding experiments.

Competition experiments

Competition experiments were largely as described (Long *et al*, 2001). For each mutant, we chose a protein concentration that resulted in 80–90% of the labeled RNA forming an RNP. For each reaction, 1.5 fmol of labeled pre-tRNA was mixed with unlabeled

competitor RNA (3.75 fmol to 15 pmol) in the presence of truncated wild-type or mutant *T. brucei* La proteins. Reaction conditions and gel analyses were as described above. Competitor RNAs consisted of either the identical RNA (containing 3' UUU_{OH}) or the RNA with a 3' UUU_p terminus. To prepare pre-tRNA^{Arg}_{CCG} with a 3' UUU_p, a pre-tRNA containing four 3' uridylates was first synthesized using T7 RNA polymerase. This was converted to the pre-RNA with a 3' UUU_p by incubating the RNA in 100 µl 1 M DL-lysine-Cl and 0.025 M NaIO₄ (pH 8.3) for 2 h at 45°C, followed by phenol extraction, ethanol precipitation, and gel purification. This Whitfield degradation reaction removes the last nucleotide, leaving a 3' phosphate (Neu and Heppel, 1964). Data were quantitated using a PhosphorImager and analyzed as described (Long *et al*, 2001).

Acknowledgements

We thank Elisabetta Ullu for her gift of the *T. brucei* La cDNA. We are grateful to Anand Saxena and the staff at X12-C at Brookhaven National Laboratory for help with data collection. We thank David W Rodgers and Elisabetta Ullu for critically reading and discussing this manuscript and the reviewers for their comments. This work was supported by grants from the Pew Foundation and the Mathers Foundation to KMR and grant R01-GM48410 from the National Institute of Health to SLW. SLW is an Associate Investigator of the Howard Hughes Medical Institute.

References

- Aigner S, Postberg J, Lipps HJ, Cech TR (2003) The Euplotes La motif protein p43 has properties of a telomerase-specific subunit. *Biochem J* 42: 5736–5747
- Allers J, Shamoo Y (2001) Structure-based analysis of protein–RNA interactions using the program ENTANGLE. *J Mol Biol* 311: 75–86
- Alspaugh MA, Tan EM (1975) Antibodies to cellular antigens in Sjogren's syndrome. *J Clin Invest* 55: 1067–1073
- Antson A (2000) Single stranded RNA binding proteins. *Curr Opin Struct Biol* 10: 87–94
- Bai C, Li Z, Tolias PP (1994) Developmental characterization of a *Drosophila* RNA-binding protein homologous to the human systemic lupus erythematosus-associated La/SS-B autoantigen. *Mol Biol Cell* 14: 5123–5129
- Brunger AT, Adams PD, Clore GM, DeLano WL, Gros P, Grosse-Kunstleve RW, Jiang JS, Kuszewski J, Nilges M, Pannu NS, Read RJ, Rice LM, Simonson T, Warren GL (1998) Crystallography and NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr D* 54: 905–921
- Chakshumathi G, Kim SD, Rubinson DA, Wolin SL (2003) A La protein requirement for pre-tRNA folding. *EMBO J* 22: 6562–6572
- Chambers JC, Kenan DJ, Martin BJ, Keene JD (1988) Genomic structure and amino acid sequence domains of the human La autoantigen. *J Biol Chem* 263: 18043–18051
- Chong SS, Hu P, Hernandez N (2001) Reconstitution of transcription from the human U6 small nuclear RNA promoter with eight recombinant polypeptides and a partially purified RNA polymerase III complex. *J Biol Chem* 276: 20727–20734
- Crowder SM, Kanaar R, Rio DC, Alber T (1999) Absence of inter-domain contacts in the crystal structure of the RNA recognition motifs of Sex-lethal. *Proc Natl Acad Sci USA* 96: 4892–4897
- Double S (1997) Preparation of selenomethionyl proteins for phase determination. In *Macromolecular Crystallography*, CW Carter, RM Sweet (eds) Vol. 276, pp 523–529. New York: Academic Press
- Fan H, Sakulich AL, Goodier JL, Zhang X, Qin J, Marais RJ (1997) Phosphorylation of the human La protein on serine 366 can regulate recycling of RNA polymerase III transcription complexes. *Cell* 88: 707–715
- Fan H, Goodier JL, Chamberlain JR, Engelke DR, Marais RJ (1998) 5' processing of tRNA precursors can be modulated by the human La antigen phosphoprotein. *Mol Cell Biol* 18: 3201–3211
- Fourmy D, Guittet E, Yoshizawa S (2002) Structure of prokaryotic SECIS mRNA hairpin and its interaction with elongation factor SelB. *J Mol Biol* 324: 137–150
- Gajiwala KS, Burley SK (2000) Winged helix proteins. *Curr Opin Struct Biol* 10: 110–116
- Goodier JL, Fan H, Marais RJ (1997) A carboxy-terminal basic region controls RNA polymerase III transcription factor activity of human La protein. *Mol Cell Biol* 17: 5823–5832
- Gottlieb E, Steitz JA (1989) The RNA binding protein La influences both the accuracy and the efficiency of the RNA polymerase III transcription *in vitro*. *EMBO J* 8: 841–850
- Jacks A, Babon J, Kelly G, Manolaridis I, Cary PD, Curry S, Conte MR (2003) Structure of the C-terminal domain of human La protein reveals a novel RNA recognition motif coupled to a helical nuclear retention element. *Structure* 11: 833–843
- Kleywegt G, Jones AT (1997) Electron density map interpretation. In *Methods in Crystallography*, CW Carter, RM Sweet (eds) Vol. 277, pp 173–208. New York: Academic Press
- Kraulis PJ (1991) MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures. *J Appl Crystallogr* 24: 946–950
- Kufel J, Allmang C, Chanfreau G, Petfalski E, Lafontaine DL, Tollervey D (2000) Precursors to the U3 small nucleolar RNA lack small nucleolar RNP proteins but are stabilized by La binding. *Mol Cell Biol* 20: 5415–5424
- Lin-Marq N, Clarkson SG (1998) Efficient synthesis, termination, and release of RNA polymerase III transcripts in *Xenopus* extracts depleted of La protein. *EMBO J* 17: 2033–2041
- Long SK, Cedervall T, Walch-Solimena C, Noe DA, Huddleston MJ, Annan RS, Wolin SL (2001) Phosphorylation of the *Saccharomyces cerevisiae* La protein does not appear to be required for its function in tRNA maturation and nascent RNA stabilization. *RNA* 7: 1589–1602
- Marais RJ (1996) Transcription termination factor La is also an initiation factor for RNA polymerase III. *Proc Natl Acad Sci USA* 93: 3383–3387
- Marais RJ, Intine RV (2001) Recognition of nascent RNA by the human La antigen: conserved and divergent features of structure and function. *Mol Cell Biol* 21: 367–379
- Marchetti MA, Tschudi C, Kwon H, Wolin SL, Ullu E (2000) Import of proteins into the trypanosome nucleus and their distribution at karyogenesis. *J Cell Sci* 113: 899–906
- Mattioli M, Reichlin M (1974) Heterogeneity of RNA protein antigens reactive with sera of patients with systemic lupus erythematosus. Descriptions of a cytoplasmic nonribosomal antigen. *Arthritis Rheum* 17: 421–429
- Mer G, Bochkareva A, Gupta R, Bochkareva E, Frappier L, Ingles CJ, Edwards AM, Chazin WJ (2000) Structural basis for the recognition of DNA repair proteins UNG2, XPA, and RAD52 by replication factor RPA. *Cell* 103: 449–456

- Neu HC, Heppel LA (1964) Nucleotide sequence analysis of polyribonucleotides by means of periodate oxidation followed by cleavage with an amine. *J Biol Chem* **237**: 2927–2934
- Nicholls A, Sharp KA, Honig B (1991) Protein folding and association: insights from the interfacial and thermodynamic properties of hydrocarbons. *Proteins* **11**: 281–296
- Ohndorf U-M, Steegborn C, Knijff R, Sondermann P (2001) Contribution of the individual domains in human La protein and its RNA 3' end-binding activity. *J Biol Chem* **276**: 27188–27196
- Otwinowski Z, Minor W (1997) Processing X-ray diffraction data collected in oscillation mode. In *Macromolecular Crystallography*, CW Carter, RM Sweet (eds) Vol. 276, pp 3–26. New York: Academic Press
- Pannone BK, Xue D, Wolin SL (1998) A role for the yeast La protein in U6 snRNP assembly: evidence that the La protein is a molecular chaperone for RNA polymerase III transcripts. *EMBO J* **17**: 7442–7453
- Rodgers D (1997) Practical cryocrystallography. In *Methods in Enzymology*, CW Carter and RM Sweet (eds) Vol. 276, pp 183–202. New York: Academic Press
- Selmer M, Su X-D (2002) Crystal structure of an mRNA-binding fragment of *Moorella thermoacetica* elongation factor SelB. *EMBO J* **21**: 4145–4155
- Sobel SG, Wolin SL (1999) Two yeast La motif-containing proteins are RNA-binding proteins that associate with polyribosomes. *Mol Biol Cell* **10**: 3849–3862
- Stefano JE (1984) Purified lupus antigen La recognizes an oligouridylylate stretch common to the 3' termini of RNA polymerase III transcripts. *Cell* **36**: 145–154
- Van Horn DJ, Yoo CJ, Xue D, Shi H, Wolin SL (1997) The La protein in *Schizosaccharomyces pombe*: a conserved yet dispensable phosphoprotein that functions in tRNA maturation. *RNA* **3**: 1434–1443
- Varani G, Nagai K (1998) RNA recognition by RNP proteins during RNA processing. *Annu Rev Biophys Biomol Struct* **27**: 407–445
- Wah DA, Hirsch JA, Dorner LF, Schildkraut I, Aggarwal AK (1997) Structure of the multimodular endonuclease *FokI* bound to DNA. *Nature* **388**: 97–100
- Wolin SL, Cedervall T (2002) The La protein. *Annu Rev Biochem* **71**: 375–403
- Wuttke DS, Foster MP, Case DA, Gottesfeld JM, Wright PE (1997) Solution structure of the first three Zinc fingers of TFIIIA bound to the cognate DNA sequence: determinants of affinity and sequence specificity. *J Mol Biol* **273**: 183–206
- Xue D, Rubinson DA, Pannone BK, Yoo CJ, Wolin SL (2000) U6 snRNP assembly in yeast involves the La protein. *EMBO J* **19**: 1650–1660
- Yisraeli JK, Melton DA (1989) Synthesis of long, capped transcripts *in vitro* by SP6 and T7 RNA polymerases. *Methods Enzymol* **180**: 42–50
- Yoo CJ, Wolin SL (1994) La proteins from *Drosophila melanogaster* and *Saccharomyces cerevisiae*: a yeast homolog of the La autoantigen is dispensable for growth. *Mol Cell Biol* **14**: 5412–5424
- Yoo CJ, Wolin SL (1997) The yeast La protein is required for the 3' endonucleolytic cleavage that matures tRNA precursors. *Cell* **89**: 393–402